

Regularity of harmonic functions for anisotropic fractional Laplacian

Paweł Sztonyk

June 4, 2007

Abstract

We prove that bounded harmonic functions of anisotropic fractional Laplacians are Hölder continuous under mild regularity assumptions on the corresponding Lévy measure. Under some stronger assumptions the Green function, Poisson kernel and the harmonic functions are even differentiable of order up to three.

1 Introduction

Let $d \geq 2$ be a natural number and let $\alpha \in (0, 2)$. We investigate a convolution semigroup of nondegenerate symmetric α -stable probability measures $\{P_t, t > 0\}$ on $\mathbb{R}^d$ with the corresponding Lévy measure ν (for definitions see Section 2). Each P_t has a smooth density p_t . We consider the *potential measure* $\mathbb{V} = \int_0^\infty P_t dt$ and the *potential kernel*

$$V(x) = \int_0^\infty p_t(x) dt, \quad x \in \mathbb{R}^d.$$

We recall that $V(x) = |x|^{\alpha-d}V(x/|x|)$, but the function may be infinite in some directions ([12, pp. 148-149]). It is known that if ν is a γ -measure on $\mathbb{S}$

⁰ Institute of Mathematics and Computer Science, Wrocław University of Technology, Wybrzeże Wyspiańskiego 27, 50-370 Wrocław, Poland
e-mail: sztonyk@pwr.wroc.pl

⁰2000 *MS Classification*: Primary 47D03, 31C05; Secondary 60J35, 60G51.

Key words and phrases: potential kernel, Green function, harmonic function, Hölder continuity, stable process.

Research supported by KBN 1 P03A 026 29 and The Alexander von Humboldt Foundation

(see (4)) with $\gamma > d - 2\alpha$, then V is continuous hence bounded on $\mathbb{S}$ (see [13, Theorem 1]). Conversely, if V is bounded on $\mathbb{S}$ then ν is a $(d - 2\alpha)$ - measure on $\mathbb{S}$ ([13, Theorem 2]).

In the present paper we study Hölder continuity and differentiability of V . In what follows we assume that ν is a γ -measure on $\mathbb{S}$, see (4). We denote

$$\kappa_0 = \gamma - (d - 2\alpha).$$

We note that $\gamma \leq d$, therefore $\kappa_0 \leq 2\alpha < 4$. The following theorem is our main result on the regularity of the potential kernel (for the definition of the relevant function spaces see below).

THEOREM 1 *Assume that $\kappa_0 > 0$. If $\kappa_0 \notin \mathbb{N}$ then the potential kernel $V(x)$ belongs to the local Hölder space $C_{loc}^{\kappa_0}(\mathbb{R}^d \setminus \{0\})$. If $\kappa_0 \in \mathbb{N}$ then $V(x) \in C_{loc}^{\kappa_0-}(\mathbb{R}^d \setminus \{0\})$. For every multiindex β such that $0 \leq |\beta| < \kappa_0$ there exists a constant $c = c(\beta)$ such that*

$$(1) \quad |D^\beta V(x)| \leq c|x|^{\alpha-d-|\beta|}, \quad |x| > 0.$$

We define operator $\mathcal{A}$ on smooth functions φ with compact support in $\mathbb{R}^d$, $\varphi \in C_c^\infty(\mathbb{R}^d)$, by letting

$$\begin{aligned} \mathcal{A}\varphi(x) &= \int_{\mathbb{R}^d} (\varphi(x+y) - \varphi(x) - y \cdot \nabla \varphi(x) \mathbf{1}_{|y|<1}) \nu(dy) \\ &= \lim_{\varepsilon \rightarrow 0^+} \int_{|y|>\varepsilon} (\varphi(x+y) - \varphi(x)) \nu(dy). \end{aligned}$$

$\mathcal{A}$ is called anisotropic fractional Laplacian and it is a restriction of the infinitesimal generator of $\{P_t\}$ [24, Example 4.1.12]. In the special case of $\nu(dy) = c|y|^{-d-\alpha}dy$ we obtain the fractional Laplacian $\Delta^{\alpha/2}$. For properties of $\Delta^{\alpha/2}$ and a discussion of equivalent definitions of its harmonic functions we refer the reader to [5, 9].

Harmonic functions corresponding to $\mathcal{A}$, or ν , are defined by the mean value property with respect to an appropriate family of harmonic measures, see Section 2. We denote

$$\kappa_1 = \gamma - (d - \alpha),$$

and we observe that $\kappa_1 = \kappa_0 - \alpha$ and $\kappa_1 \leq \alpha < 2$.

THEOREM 2 *If $\kappa_1 > 0$ then there exists a constant $\rho \in (0, \kappa_1]$ such that every function u bounded on $\mathbb{R}^d$ and harmonic in $B(0, 1)$ belongs to the local Hölder space $C_{loc}^\rho(B(0, 1))$ and*

$$(2) \quad \|u\|_{C^\rho(B(0, 1-r))} \leq cr^{-\rho}\|u\|_\infty, \quad r \in (0, 1/2].$$

If $\kappa_1 \leq \alpha < 1$ then we can take $\rho = \kappa_1$ and if $\kappa_1 > 1$ then $\rho \in (1, \kappa_1)$.

In particular every bounded function harmonic in $B(0, 1)$ is Hölder continuous if $\kappa_1 \in (0, 1]$ and even differentiable if $\kappa_1 > 1$. We note that for $\alpha \geq 1$ an explicit value of ρ can be obtained from Lemma 11 and (46).

Here are a few comments about the general context of our study. A primary source of motivation for this study is the potential theory of the second order elliptic partial differential operators. Hölder continuity of harmonic functions of the second order elliptic partial differential operators in divergence form was proved independently in [18] and [35]. J. Moser in [34] gave another method of the proof using Harnack's inequality and an iteration technique that has been later used successfully in many other situations. In these papers the local character of the operators is crucial. Perturbations of strongly elliptic operators by 'smaller' nonlocal operators can also be dealt with as shown in [32, 21, 27].

Harnack's inequality and Hölder continuity for bounded harmonic functions of purely nonlocal integro-differential operators with kernels $\nu(x, dy)$ were obtained in [2]. We note that the results of [2] were restricted to absolutely continuous $\nu(x, dy)$ with densities comparable to $|y - x|^{-d-\alpha}$. The class of considered operators then gradually extended, see [40] and [39]. In [1] the Hölder continuity is proved without the assumption that the density exists. Instead, the measures $\nu(x, \cdot)$ considered in [1] satisfy some regularity conditions and estimates from below. Similar, slightly weaker assumptions yield in [22] logarithmic modulus of continuity for harmonic functions: $|u(y) - u(x)| \leq c\|u\|_\infty |\log |y - x||^{-\rho}$.

In our study, initiated in [10] (see also [11, 12, 13]), we restrict ourselves to the translation invariant case $\nu(x, A) = \nu(0, A - x) = \nu(A - x)$ but we do not generally rely on the existence of the density of the Lévy measure ν , and we do not assume estimates for ν from below. We thus allow for much more anisotropy than [2, 40, 39, 1, 22]. It should be made clear that while our study is analogous to the study of symmetric second order elliptic differential operators with mere constant coefficients the variety of different behaviours of the resulting fractional Laplacians makes the subject very complex and

interesting. We believe that the study paves a way to further generalizations in spirit of [2] and may also contribute to the general potential theory.

The main focus of the present paper are estimates of the potential kernel $V(x)$ (Theorem 1). This is a prominent harmonic function of $\mathcal{A}$ on $\mathbb{R}^d \setminus \{0\}$, see [12], and results for more general harmonic functions (Theorem 2) may be considered as consequences of such estimates. For sufficiently regular ν and $\alpha > 1/2$ we obtain the differentiability of harmonic functions which is a new result even for absolutely continuous measures. Our methods are mainly analytic although we use a probabilistic framework to define some of the objects in question. We do not use Moser's method here because the Harnack's inequality may not hold even if ν has a bounded density (see [12]). We base our development on certain estimates of the derivatives of transition densities (Lemma 3), extending some of the results of [29, 13, 44, 36, 20].

Apart from Theorem 1 and 2 we obtain here several results concerning regularity of transition densities, Green function and Poisson kernel for the ball (see Lemma 3, Proposition 1, Lemma 9, 10 and 17). These objects were investigated in detail in the isotropic case $\nu(dy) = c|y|^{-d-\alpha}dy$. The results obtained include estimates of the Green function and Poisson kernel (see [28, 26, 31, 30, 6]), estimates of the derivatives of transition density and resolvent (see [7, 29]) and gradient estimates of harmonic functions (see [8]). The main difficulty of the present study in comparison to the isotropic case is the lack of explicit formulas, e.g. for the Poisson kernel or the Green function for a ball.

The paper is organized as follows. Definitions and preliminaries are given in Section 2. In Section 3 we estimate the derivatives of the transition densities extending the estimates of the semigroup and potential kernel given in [13]. At the end of Section 3 we prove Theorem 1. The continuity estimates of the Green function and its derivative are proved in Section 4. In Section 5 we investigate the regularity of the Poisson kernel. Theorem 2 is proved in Section 6.

2 Preliminaries

The general references for this section are [14, 15, 16, 38, 3, 4]. Recall that $d \geq 2$ and $\alpha \in (0, 2)$. We explicitly exclude $d = 1$ as it leads to the well-know rotation invariant case (for which see, e.g., [5, 9]).

For $x \in \mathbb{R}^d$ and $r > 0$ we let $|x| = \sqrt{\sum_{i=1}^d x_i^2}$ and $B(x, r) = \{y \in \mathbb{R}^d :$

$|y - x| < r\}$. We denote $\mathbb{S} = \{x \in \mathbb{R}^d : |x| = 1\}$. All the sets, functions and measures considered in the sequel will be Borel. For a measure λ on $\mathbb{R}^d$, $|\lambda|$ denotes its total mass. We call λ degenerate if there is a proper linear subspace M of $\mathbb{R}^d$ such that $\text{supp}(\lambda) \subset M$; otherwise we call λ *nondegenerate*.

In what follows we will consider measures μ concentrated on $\mathbb{S}$. We will assume that μ is positive, finite, nondegenerate (in particular $\mu \neq 0$), and symmetric:

$$\mu(D) = \mu(-D), \quad D \subset \mathbb{S}.$$

We will call μ the *spectral measure*. We let

$$(3) \quad \nu(D) = \int_{\mathbb{S}} \int_0^\infty \mathbf{1}_D(r\xi) r^{-1-\alpha} dr \mu(d\xi), \quad D \subset \mathbb{R}^d,$$

where $\mathbf{1}_D$ is the indicator function of D . Note that ν is symmetric. It is a Lévy measure on $\mathbb{R}^d$, i.e.

$$\int_{\mathbb{R}^d} \min(|y|^2, 1) \nu(dy) < \infty.$$

For $r > 0$ and a function φ on $\mathbb{R}^d$ we consider its dilation $\varphi_r(y) = \varphi(y/r)$, and we note that $\nu(\varphi_r) = r^{-\alpha} \nu(\varphi)$. In particular ν is homogeneous: $\nu(rB) = r^{-\alpha} \nu(B)$ for $B \subset \mathbb{R}^d$.

We consider, after [13], an auxiliary scale of smoothness for ν .

DEFINITION 1 *We say that ν is a γ -measure on $\mathbb{S}$ if there exists a constant c such that*

$$(4) \quad \nu(B(x, r)) \leq cr^\gamma, \quad |x| = 1, \quad 0 < r < 1/2.$$

Since $\nu(dr d\theta) = r^{-1-\alpha} dr \mu(d\theta)$, it is at least a 1-measure and at most a d -measure on $\mathbb{S}$. If ν is a γ -measure with $\gamma > 1$, then μ has no atoms. Finally, ν is a d -measure if and only if it is absolutely continuous with respect to the Lebesgue measure and its density function is locally bounded on $\mathbb{R}^d \setminus \{0\}$.

We consider the semigroup of *stable* probability measures $\{P_t, t > 0\}$ with the Fourier transform $\mathcal{F}(P_t)(u) = \exp(-t\Phi(u))$, where

$$\begin{aligned} \Phi(u) &= - \int (e^{iu \cdot y} - 1 - iu \cdot y \mathbf{1}_{B(0,1)}(y)) \nu(dy) \\ &= - \int (\cos(u \cdot y) - 1) \nu(dy) = \frac{\pi}{2 \sin \frac{\pi\alpha}{2} \Gamma(1 + \alpha)} \int_{\mathbb{S}} |u \cdot \xi|^\alpha \mu(d\xi). \end{aligned}$$

Here $u \cdot \xi$ denotes the usual inner product of $u, \xi \in \mathbb{R}^d$. Since μ is finite, homogeneous and nondegenerate,

$$(5) \quad \Phi(u) = |u|^\alpha \Phi(u/|u|) \approx |u|^\alpha.$$

We also let $\bar{\nu} = \mathbf{1}_{B(0,1)^c} \nu$, and $\tilde{\nu} = \mathbf{1}_{B(0,1)} \nu$ and consider the corresponding semigroups of measures $\{\tilde{P}_t, t \geq 0\}$ and $\{\bar{P}_t, t \geq 0\}$ such that

$$\begin{aligned} \mathcal{F}(\bar{P}_t)(u) &= \int e^{iuy} \bar{P}_t(dy) = \exp \left(t \int (\cos(u \cdot y) - 1) \bar{\nu}(dy) \right), \quad u \in \mathbb{R}^d, \\ (6) \quad \mathcal{F}(\tilde{P}_t)(u) &= \exp \left(t \int (\cos(u \cdot y) - 1) \tilde{\nu}(dy) \right), \quad u \in \mathbb{R}^d. \end{aligned}$$

We then have that

$$P_t = \tilde{P}_t * \bar{P}_t, \quad t \geq 0.$$

The measures P_t and $\tilde{P}_t$ have rapidly decreasing Fourier transform hence they are absolutely continuous with smooth bounded densities denoted $p_t(x)$ and $\tilde{p}_t(x)$, respectively. Of course,

$$(7) \quad p_t = \tilde{p}_t * \bar{P}_t, \quad t > 0.$$

By using (5) we obtain the *scaling property* of $\{p_t\}$:

$$(8) \quad p_t(x) = k^d p_{k^\alpha t}(kx), \quad x \in \mathbb{R}^d, k > 0,$$

or

$$p_t(x) = t^{-d/\alpha} p_1(t^{-1/\alpha} x), \quad x \in \mathbb{R}^d.$$

In particular,

$$(9) \quad p_t(x) \leq ct^{-d/\alpha}.$$

If ν is a γ - measure on $\mathbb{S}$ then we have

$$(10) \quad p_1(x) \leq c(1 + |x|)^{-\alpha-\gamma},$$

see [13] or [44]. We define the *potential measure* of the semigroup $\{P_t\}$:

$$\mathbb{V}(D) = \int_0^\infty P_t(D) dt, \quad D \subset \mathbb{R}^d.$$

Since $\alpha < d$, by (9), $\mathbb{V}$ is finite on bounded subsets of $\mathbb{R}^d$. Let

$$(11) \quad V(x) = \int_0^\infty p_t(x) dt, \quad x \in \mathbb{R}^d,$$

so that

$$\mathbb{V}(D) = \int_D V(x) dx, \quad D \subset \mathbb{R}^d.$$

We call $V(x)$ the *potential kernel* of the stable semigroup. By (8)

$$(12) \quad V(x) = |x|^{\alpha-d} V(x/|x|), \quad x \neq 0.$$

If ν is a γ -measure on $\mathbb{S}$ and $\gamma > d - 2\alpha$ then $V(x)$ is continuous on $\mathbb{S}$ ([13, Theorem 1]).

The Lévy measure ν determines the symmetric stable Lévy process (X_t, P^x) with generating triplet $(0, \nu, 0)$. The transition probabilities of the process (X_t, P^x) are $P(t, x, A) = P_t(A - x)$, $t > 0$, $x \in \mathbb{R}^d$, $A \subset \mathbb{R}^d$, and $P(0, x, A) = \mathbf{1}_A(x)$. The trajectories are right continuous with left limits and the process is strong Markov with respect to the so-called *standard filtration* (see, e.g., [38], [3], or [4]).

For open $U \subset \mathbb{R}^d$ we denote $\tau_U = \inf\{t \geq 0; X_t \notin U\}$, the *first exit time* of U . We write ω_D^x for the harmonic measure of (open) D :

$$\omega_D^x(A) = P^x(\tau_D < \infty, X_{\tau_D} \in A), \quad x \in \mathbb{R}^d, A \subset \mathbb{R}^d.$$

By the strong Markov property

$$(13) \quad \omega_D^x(A) = \int \omega_D^y(A) \omega_U^x(dy), \quad \text{if } U \subset D.$$

We say that a function u on $\mathbb{R}^d$ is *harmonic* in open $D \subset \mathbb{R}^d$ if

$$(14) \quad u(x) = E^x u(X_{\tau_U}) = \int_{U^c} u(y) \omega_U^x(dy), \quad x \in \mathbb{R}^d,$$

for every bounded open set U with the closure $\bar{U}$ contained in D . It is called *regular harmonic* in D if (14) holds for $U = D$. If D is unbounded then $E^x u(X_{\tau_D}) = E^x[\tau_D < \infty; u(X_{\tau_D})]$ by a convention. Under (14) it will be only assumed that the expectation in (14) is well defined (but not necessarily finite). Regular harmonicity implies harmonicity, and is inherited by subsets $U \subset D$. This follows from (13).

We denote by $p_t^D(x, v)$ the transition density of the process killed at the first exit from D :

$$p_t^D(x, v) = p(t, x, v) - E^x[\tau_D < t; p(t - \tau_D, X_{\tau_D}, v)], \quad t > 0, \quad x, v \in \mathbb{R}^d.$$

Here $p(t, x, v) = p_t(v - x)$. We will assume in the sequel that D is *regular*: $P^x[\inf\{t > 0 : X_t \notin D\} = 0] = 1$ for $x \in D^c$ (see [15, 14]). It is well-known that p_t^D is symmetric: $p_t^D(x, v) = p_t^D(v, x)$, $x, v \in \mathbb{R}^d$ (see, e.g., [16]). The strong Markov property yields

$$(15) \quad p(t, x, v) = E^x[p(t - \tau_D, X_{\tau_D}, v); \tau_D < t], \quad x \in D, \quad v \in D^c,$$

thus $p_t^D(x, v) = 0$ if $x \in D$, $v \in D^c$. We let

$$G_D(x, v) = \int_0^\infty p_t^D(x, v) dt,$$

and we call $G_D(x, v)$ the *Green function* for D . If V is continuous on $\mathbb{R}^d \setminus \{0\}$, so that $V(x) \leq c|x|^{\alpha-d}$, then the strong Markov property yields for $x, v \in D$

$$(16) \quad G_D(x, v) = V(x, v) - E^x V(X_{\tau_D}, v) = V(x, v) - \int_{D^c} V(z, v) \omega_D^x(dz).$$

Here $V(x, v) = V(v - x)$. This holds, e.g., if ν is a γ -measure on $\mathbb{S}$ and $\gamma > d - 2\alpha$ (see [13, Theorem 1]). In particular

$$(17) \quad G_D(x, v) \leq c|v - x|^{\alpha-d}, \quad x, v \in D.$$

The Green function is symmetric: $G_D(x, v) = G_D(v, x)$, continuous in $D \times D \setminus \{(x, v) : x = v\}$, and it vanishes if $x \in D^c$ or $v \in D^c$.

For each $v \in \mathbb{R}^d$, $V(x, v)$ is harmonic in $x \in \mathbb{R}^d \setminus \{v\}$. Indeed, if $x \in D$ and $\text{dist}(D, v) > 0$ then by (15)

$$V(x, v) = \int_0^\infty E^x[p(t - \tau_D, X_{\tau_D}, v); \tau_D < t] dt = E^x V(X_{\tau_D}, v).$$

Similarly, the Green function $x \mapsto G_D(x, v)$ is harmonic in $D \setminus \{v\}$.

We like to note that $V(x, v)$ may be infinite at some points $x \neq v$, see Introduction. Thus harmonic functions need to be defined by the mean value

property (14) rather than in terms of the anisotropic fractional Laplacian $\mathcal{A}$. However, if a function φ belongs to the domain of the infinitesimal generator $\mathcal{A}'$ of the process, which is an extension of $\mathcal{A}$, and $\mathcal{A}'\varphi(x) = 0$ for every $x \in D$ then φ is harmonic in D in the sense of (14). It follows from the fact that

$$\varphi(X_t) - \varphi(x) - \int_0^t \mathcal{A}'\varphi(X_s)ds$$

is a martingale with respect to P^x (see, e.g., [17]).

By Ikeda–Watanabe formula [23] we have

$$(18) \quad \omega_D^x(A) = \int_D G_D(x, v) \nu(A - v) dv, \quad \text{if } \text{dist}(A, D) > 0.$$

If the boundary of D is smooth or even Lipschitz then

$$\omega_D^x(\partial D) = 0, \quad x \in D,$$

see ([41], [33], [45]). In this case ω_D^x is absolutely continuous with respect to the Lebesgue measure on D^c . Its density function, or the Poisson kernel, is given by the formula (see [13])

$$(19) \quad P_D(x, z) = \int_{z-D} G_D(x, z - v) \nu(dv), \quad x \in D.$$

Note that such D is regular, which follows from scaling and the fact that

$$(20) \quad p_t(x) > 0, \quad x \in \mathbb{R}^d \quad (t > 0),$$

(see [42] or [36, Lemma 5]). In particular the above considerations apply to $D = B(0, 1)$.

It follows from (5) that for every $r > 0$ and $x \in \mathbb{R}^d$ the P^x distribution of $\{X_t, t \geq 0\}$ is the same as the P^{rx} distribution of $\{r^{-1}X_{r^\alpha t}, t \geq 0\}$. In consequence,

$$(21) \quad \omega_D^x(A) = \omega_{rD}^{rx}(rA),$$

which we will call scaling, too. It yields that for u harmonic on D , the dilation, u_r , is harmonic on rD . A similar remark applies to translations.

Let $\mathbb{N}_0 = \{0, 1, 2, \dots\}$. For a multiindex $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{N}_0^d$ we denote $|\beta| = \beta_1 + \dots + \beta_d$. Let $s \geq 0$ and $s = [s] + \{s\}$ where $[s] \in \mathbb{Z}$ and $\{s\} \in [0, 1)$.

For every $s > 0$, $s \notin \mathbb{N}$ and open set $D \subset \mathbb{R}^d$ we denote by $C^s(D)$ the Hölder space of order s (see, e.g., [43]), i.e.,

$$C^s(D) = \{f \in C^{[s]}(D) : \|f\|_{C^s(D)} < \infty\}$$

where $C^0(D)$ denotes the set of bounded and uniformly continuous functions on D and for $n \in \mathbb{N}$, $C^n(D) = \{f : D^\beta f \in C^0(D) \text{ for all } |\beta| \leq n\}$, $\|f\|_{C^n(D)} = \sum_{|\beta| \leq n} \|\mathbf{1}_D \cdot D^\beta f\|_\infty$, and

$$\|f\|_{C^s(D)} = \|f\|_{C^{[s]}(D)} + \sum_{|\beta|=[s]} \sup_{x,y \in D} \frac{|D^\beta f(y) - D^\beta f(x)|}{|y - x|^{\{s\}}}.$$

We write $f \in C^{s-}(D)$ if $f \in C^{s-\delta}(D)$ for every $\delta \in (0, s]$. By $C_{loc}^s(D)$ we denote the local Hölder space, i.e. a function f belongs to $C_{loc}^s(D)$ if $f \in C^s(U)$ for every bounded open set U with the closure $\bar{U}$ contained in D .

We use c (with subscripts) to denote finite positive constants which depend only on the measure μ and the constant γ , the dimension d and the index α . Any *additional* dependence is explicitly indicated by writing, e.g., $c = c(\beta, n)$. The value of c , when used without subscripts, may change from place to place.

3 Transition density and potential kernel

We first investigate the regularity of the density of the semigroup $\{\tilde{P}_t, t \geq 0\}$ generated by the measure $\tilde{\nu} = \mathbf{1}_{B(0,1)}\nu$. Let $S(\mathbb{R}^d)$ denote the usual Schwarz space of smooth rapidly decreasing functions (see, e.g., [37]).

LEMMA 1 $\mathcal{F}(\tilde{P}_t) \in S(\mathbb{R}^d)$.

Proof. Let $\varphi(u, y) = \cos(u \cdot y) - 1$, and $\tilde{\Phi}(u) = \int \varphi(u, y) \tilde{\nu}(dy)$, see (6), where $u = (u_1, \dots, u_d)$, $y = (y_1, \dots, y_d)$. For every $\beta = (\beta_1, \dots, \beta_d) \in \mathbb{N}_0^d$, $|\beta| > 0$, we have

$$D_u^\beta \varphi(u, y) = y^\beta \cos^{(|\beta|)}(u \cdot y)$$

where $y^\beta = y_1^{\beta_1} \dots y_d^{\beta_d}$ and $\cos^{(n)}(\theta)$ denotes the n -th derivative of the function $\mathbb{R} \ni \theta \rightarrow \cos(\theta)$. Hence

$$|D_u^\beta \varphi(u, y)| \leq |y|^{|\beta|+1} (|u| \wedge \frac{1}{|y|}), \quad |\beta| = 1, 3, 5, \dots,$$

and

$$|D_u^\beta \varphi(u, y)| \leq |y|^{|\beta|}, \quad |\beta| = 2, 4, 6, \dots$$

Therefore, we can change the order of differentiation and integration, obtaining

$$D^\beta \tilde{\Phi}(u) = \int D_u^\beta \varphi(u, y) \tilde{\nu}(dy).$$

Hence

$$|D^\beta \tilde{\Phi}(u)| \leq c(\beta)g(u), \quad u \in \mathbb{R}^d, \quad |\beta| = 1, 2, 3, 4, \dots,$$

where

$$g(u) = \begin{cases} 1 & \text{for } |\beta| = 2, 4, 6, \dots, \\ |u| \wedge 1 & \text{for } |\beta| = 1, 3, 5, \dots \text{ and } |\beta| > \alpha, \\ |u| \wedge |u|^{\alpha-1} & \text{for } |\beta| = 1 \text{ and } \alpha > 1, \\ |u| \wedge (1 + |\log |u||) & \text{for } |\beta| = \alpha = 1. \end{cases}$$

We use the multivariable version of Faa di Bruno's formula (see [19]) for the function $f(u) = \mathcal{F}(\tilde{P}_t)(u) = \exp(t\tilde{\Phi}(u))$ to obtain for every $k \in \mathbb{N}$

$$\frac{\partial^k}{\partial u_1 \dots \partial u_k} f(u) = f(u) \sum_{\pi \in \mathcal{P}} \prod_{A \in \pi} \frac{\partial^{|A|}(t\tilde{\Phi})}{\prod_{j \in A} \partial u_j}(u),$$

where $\mathcal{P}$ denotes the set of all partitions of the set $\{1, \dots, k\}$. The formula is still valid if some of the variables $u_1, \dots, u_k$ on the left hand side denote the same variable and then the corresponding terms in the sum are multiply counted. Since $g(u) \leq 1 + |u|$, we get

$$|D^\beta f(u)| \leq c(\beta, t)(1 + |u|)^{|\beta|} |f(u)| \leq c(\beta, t)(1 + |u|)^{|\beta|} e^{-c|u|^\alpha}$$

□

Lemma 1 yields that for every $\beta \in \mathbb{N}_0^d$ and $q > 0$ there exists a constant $c = c(\beta, q)$ such that

$$(22) \quad |D^\beta \tilde{p}_1(y)| \leq c(1 + |y|)^{-q}, \quad y \in \mathbb{R}^d.$$

The proof of the following lemma is based on the ideas of the proofs of [36, Theorem 3] and [13, Lemma 6]. In what follows we use the convention $D^0 = \text{Id}$.

LEMMA 2 *If ν is a γ -measure then for every $\beta \in \mathbb{N}_0^d$ there exists a constant $c = c(\beta)$ such that*

$$|D^\beta p_1(y)| \leq c(1 + |y|)^{-\alpha-\gamma}, \quad y \in \mathbb{R}^d.$$

Proof. We have

$$p_1(y) = \int \tilde{p}_1(y - z) \bar{P}_1(dz).$$

Also,

$$D^\beta p_1(y) = \int D^\beta \tilde{p}_1(y - z) \bar{P}_1(dz),$$

because $\bar{P}_1$ is finite and $D^\beta \tilde{p}_1$ is bounded and continuous. By [13, Corollary 4] we have

$$\bar{P}_1(B(y, \lambda)) \leq c\lambda^\gamma(1 + \lambda^\alpha)|y|^{-\alpha-\gamma}, \quad y \in \mathbb{R}^d, \lambda > 0.$$

Hence for $q = \gamma + \alpha + 1$ by (22) we obtain

$$\begin{aligned} |D^\beta p_1(y)| &= \left| \int_{\mathbb{R}^d} D^\beta \tilde{p}_1(y - z) \bar{P}_1(dz) \right| \\ &\leq \int_{\mathbb{R}^d} |D^\beta \tilde{p}_1(y - z)| \bar{P}_1(dz) \\ &\leq \int_{\mathbb{R}^d} c(\beta)(1 + |y - z|)^{-q} \bar{P}_1(dz) \\ &= c(\beta) \int_0^1 \bar{P}_1(\{z : (1 + |z - y|)^{-q} > s\}) ds \\ &\leq c(\beta) \int_0^1 \bar{P}_1(B(y, s^{-1/q})) ds \\ &\leq c(\beta) \int_0^1 (s^{-1/q})^\gamma (1 + (s^{-1/q})^\alpha) |y|^{-\gamma-\alpha} ds \\ &= c(\beta) |y|^{-\gamma-\alpha} \left[\int_0^1 s^{-\gamma/q} ds + \int_0^1 s^{-(\gamma+\alpha)/q} ds \right] \\ &= c(\beta) |y|^{-\gamma-\alpha}. \end{aligned}$$

Since $D^\beta p_1$ is bounded we also have $|D^\beta p_1(y)| \leq c(\beta)(1 + |y|)^{-\alpha-\gamma}$. $\square$

We note that explicit estimates for the first derivative of the transition density in the isotropic case of $\nu(dy) = c|y|^{-d-\alpha}dy$ are also given in [7, Lemma 5]. In this case we have $|\nabla_y p(1, y)| \leq c|y|(1 + |y|)^{-d-2-\alpha}$.

LEMMA 3 For every $\beta \in \mathbb{N}_0^d$ there exists a constant $c = c(\beta)$ such that for every $R > 0$ we have

$$(23) \quad |D^\beta p_1(y) - D^\beta p_1(x)| \leq c(1+R)^{-\alpha-\gamma}(|y-x| \wedge 1),$$

provided $x, y \in B(0, R)^c$, $|y-x| < R/4$.

Proof. By the Lagrange's theorem we obtain

$$D^\beta p_1(y) - D^\beta p_1(x) = \nabla D^\beta p_1(x + \theta(y-x)) \cdot (y-x),$$

where $\theta \in [0, 1]$. If $|y-x| < 1$ we get by Lemma 2

$$\begin{aligned} |D^\beta p_1(y) - D^\beta p_1(x)| &\leq |\nabla D^\beta p_1(x + \theta(y-x))| \cdot |y-x| \\ &\leq c(1+|x + \theta(y-x)|)^{-\alpha-\gamma}|y-x| \\ &\leq c(1+(3/4)R)^{-\alpha-\gamma}|y-x| \\ &\leq c(1+R)^{-\alpha-\gamma}|y-x|. \end{aligned}$$

For $|y-x| \geq 1$ we have

$$|D^\beta p_1(y) - D^\beta p_1(x)| \leq 2 \sup_{z \in B(0, R)^c} |D^\beta p_1(z)| \leq c(1+R)^{-\alpha-\gamma}.$$

Proof of Theorem 1. We will first prove that for every $\beta \in \mathbb{N}_0^d$ such that $|\beta| < \kappa_0 = \gamma - (d - 2\alpha)$, we have □

$$(24) \quad D^\beta V(x) = \int_0^\infty D_x^\beta p_t(x) dt, \quad |x| > 0.$$

By scaling property (8) we have for every $\beta \in \mathbb{N}_0^d$

$$D_x^\beta p_t(x) = k^{d+|\beta|} (D_x^\beta p_{k^\alpha t})(kx), \quad x \in \mathbb{R}^d, \quad t > 0, \quad k > 0.$$

For $k = t^{-1/\alpha}$ we get by Lemma 2

$$\begin{aligned} |D_x^\beta p_t(x)| &= t^{\frac{-d-|\beta|}{\alpha}} |(D^\beta p_1)(t^{-1/\alpha}x)| \\ &\leq c(\beta) t^{\frac{-d-|\beta|}{\alpha}} (1 + t^{-1/\alpha}|x|)^{-\alpha-\gamma}, \quad x \in \mathbb{R}^d, \quad t > 0, \end{aligned}$$

and this yields (24) and the continuity of $D^\beta V(x)$ for $x \neq 0$. Hence we have

$$\begin{aligned}
(25) \quad D^\beta V(x) &= \int_0^\infty D_x^\beta p_t(x) dt \\
&= k^{d+|\beta|} \int_0^\infty (D_x^\beta p_{k^\alpha t})(kx) dt \\
&= k^{-\alpha+d+|\beta|} \int_0^\infty (D_x^\beta p_s)(kx) ds = k^{-\alpha+d+|\beta|} (D^\beta V)(kx),
\end{aligned}$$

where $k > 0$. This yields (1).

Let $|y - x| < 1/4$ and $|x| > 1$, $|y| > 1$. By scaling and Lemma 3

$$\begin{aligned}
|D^\beta p_t(y) - D^\beta p_t(x)| &= t^{\frac{-d-|\beta|}{\alpha}} |(D^\beta p_1)(t^{-1/\alpha}y) - (D^\beta p_1)(t^{-1/\alpha}x)| \\
&\leq t^{\frac{-d-|\beta|}{\alpha}} c(1 + t^{-1/\alpha})^{-\alpha-\gamma} (|t^{-1/\alpha}y - t^{-1/\alpha}x| \wedge 1) \\
&= ct^{\frac{-d-|\beta|-1}{\alpha}} (1 + t^{-1/\alpha})^{-\alpha-\gamma} (|y - x| \wedge t^{1/\alpha}), \quad t > 0.
\end{aligned}$$

This yields

$$\begin{aligned}
|D^\beta V(y) - D^\beta V(x)| &\leq c \int_0^\infty t^{\frac{-d-|\beta|-1}{\alpha}} (1 + t^{-1/\alpha})^{-\alpha-\gamma} (|y - x| \wedge t^{1/\alpha}) dt \\
&= c \int_0^{|y-x|^\alpha} t^{\frac{-d-|\beta|-1}{\alpha}} (1 + t^{-1/\alpha})^{-\alpha-\gamma} t^{1/\alpha} dt \\
&\quad + c \int_{|y-x|^\alpha}^\infty t^{\frac{-d-|\beta|-1}{\alpha}} (1 + t^{-1/\alpha})^{-\alpha-\gamma} |y - x| dt \\
&= I + II.
\end{aligned}$$

Let $\kappa_0 \notin \mathbb{N}$ and $|\beta| = [\kappa_0]$. We have

$$I \leq c \int_0^{|y-x|^\alpha} t^{\frac{-d-|\beta|+\alpha+\gamma}{\alpha}} dt = c|y-x|^{2\alpha+\gamma-d-|\beta|} = c|y-x|^{\{\kappa_0\}},$$

$$\begin{aligned}
II &\leq c \int_{|y-x|^\alpha}^1 t^{\frac{-d-|\beta|-1+\alpha+\gamma}{\alpha}} |y-x| dt + c \int_1^\infty t^{\frac{-d-|\beta|-1}{\alpha}} |y-x| dt \\
&\leq c(|y-x|^{\{\kappa_0\}} + |y-x|) \leq c|y-x|^{\{\kappa_0\}}.
\end{aligned}$$

If $|y - x| \geq \frac{1}{4}$ then also we can write

$$|D^\beta V(y) - D^\beta V(x)| \leq 2 \cdot \left(\frac{|y-x|}{4} \right)^{\{\kappa_0\}} \cdot \sup_{z \in B(0,1)^c} |D^\beta V(z)|.$$

Therefore, by (25) we have that for every $r > 0$ and $x, y \in B(0, r)^c$,

$$(26) \quad \begin{aligned} |D^\beta V(y) - D^\beta V(x)| &= r^{\alpha-d-|\beta|} |(D^\beta V)(y/r) - (D^\beta V)(x/r)| \\ &\leq cr^{\alpha-d-|\beta|} (|y-x|/r)^{\{\kappa_0\}}. \end{aligned}$$

Thus if $\kappa_0 \notin \mathbb{N}$ then $V \in C^{\kappa_0}(B(0, r)^c)$ for every $r > 0$, or $V \in C_{loc}^{\kappa_0}(\mathbb{R}^d \setminus \{0\})$. We now assume that $\kappa_0 \in \mathbb{N}$ and $|\beta| = \kappa_0 - 1$. Let $|x| > 1$, $|y| > 1$, $|y-x| < 1/4$. We have

$$\begin{aligned} I &= \int_0^{|y-x|^\alpha} t^{\frac{-d-|\beta|-1}{\alpha}} (1+t^{-1/\alpha})^{-\alpha-\gamma} t^{1/\alpha} dt \\ &\leq c \int_0^{|y-x|^\alpha} t^{\frac{1-\alpha}{\alpha}} dt \\ &= c|y-x|, \end{aligned}$$

and

$$\begin{aligned} II &= \int_{|y-x|^\alpha}^\infty t^{\frac{-d-|\beta|-1}{\alpha}} (1+t^{-1/\alpha})^{-\alpha-\gamma} |y-x| dt \\ &\leq c \int_{|y-x|^\alpha}^1 t^{-1} |y-x| dt + c \int_1^\infty t^{\frac{-d-|\beta|-1}{\alpha}} |y-x| dt \\ &= c|y-x| (\log(\frac{1}{|y-x|}) + 1). \end{aligned}$$

This yields

$$|D^\beta V(y) - D^\beta V(x)| \leq c|y-x| \log(2 + \frac{1}{|y-x|}).$$

If $|y-x| \geq \frac{1}{4}$ then we can write

$$|D^\beta V(y) - D^\beta V(x)| \leq 16|y-x| \log(2 + \frac{1}{|y-x|}) \sup_{z \in B(0,1)^c} |D^\beta V(z)|.$$

Thus, for every $r > 0$, $x, y \in B(0, r)^c$ by (25) we get

$$(27) \quad \begin{aligned} |D^\beta V(y) - D^\beta V(x)| &= r^{\alpha-d-|\beta|} |(D^\beta V)(y/r) - (D^\beta V)(x/r)| \\ &\leq cr^{\alpha-d-|\beta|} (|y-x|/r) \log(2 + \frac{r}{|y-x|}). \end{aligned}$$

In particular, $V(x) \in C^{\kappa_0-p}(B(0, r)^c)$ for every $p \in (0, 1)$. $\square$

We note that if the Lévy measure ν is absolutely continuous and has a locally bounded density, i.e., $\gamma = d$, then $\kappa_0 = 2\alpha$ and the potential kernel is differentiable for $\alpha > 1/2$ and 2α -Hölder continuous for $\alpha \in (0, 1/2)$.

4 Green function

In this section we consider the regularity of the Green function of the unit ball.

For every finite measure m we denote by $\mathbb{V}m(x)$ its potential at x :

$$\mathbb{V}m(x) = \int_{\mathbb{R}^d} V(x-z)m(dz).$$

In what follows for every $s \in \mathbb{R}$ we denote

$$[s]^- = \begin{cases} [s] & \text{for } s \notin \mathbb{Z}, \\ s-1 & \text{for } s \in \mathbb{Z}. \end{cases}$$

LEMMA 4 *Let m be a finite measure with support S and $\kappa_0 = \gamma - (d-2\alpha) > 0$. If $\kappa_0 \notin \mathbb{N}$ then the potential $\mathbb{V}m(x)$ belongs to the local Hölder space $C_{loc}^{\kappa_0}(S^c)$. If $\kappa_0 \in \mathbb{N}$ then $\mathbb{V}m(x) \in C_{loc}^{\kappa_0-}(S^c)$. Moreover there is c such that for every $r > 0$ and $|\beta| = [\kappa_0]^-$ we have*

$$(28) \quad |D^\beta \mathbb{V}m(y) - D^\beta \mathbb{V}m(x)| \leq cr^{\alpha-d-|\beta|} f\left(\frac{|y-x|}{r}\right) |m|,$$

provided $\text{dist}(x, S) > r$, $\text{dist}(y, S) > r$, where

$$f(\rho) = \begin{cases} \rho^{\{\kappa_0\}} & \text{for } \kappa_0 \notin \mathbb{N}, \\ \rho \log(2 + \frac{1}{\rho}) & \text{for } \kappa_0 \in \mathbb{N}. \end{cases}$$

For every $|\beta| < \kappa_0$ we have

$$(29) \quad |D^\beta \mathbb{V}m(x)| \leq c \text{dist}(x, S)^{\alpha-d-|\beta|} |m|, \quad x \in S^c.$$

Lemma 4 follows easily from Theorem 1, (26) and (27) and so we omit the proof.

In what follows we denote $B = B(0, 1)$.

PROPOSITION 1 *Let $\kappa_0 = \gamma - (d-2\alpha) > 0$. If $\kappa_0 \notin \mathbb{N}$ then for every $v \in B$ the function $g(x) = G_B(x, v)$ belongs to the local Hölder space $C_{loc}^{\kappa_0}(B \setminus \{v\})$. If $\kappa_0 \in \mathbb{N}$ then $g(x) \in C_{loc}^{\kappa_0-}(B \setminus \{v\})$. Moreover there is c such that for every $r \in (0, 1)$ and $|\beta| = [\kappa_0]^-$ we have*

$$(30) \quad |D^\beta g(y) - D^\beta g(x)| \leq cr^{\alpha-d-|\beta|} f\left(\frac{|y-x|}{r}\right),$$

provided $x, y \in B(0, 1 - r) \cap B(v, r)^c$, where

$$f(\rho) = \begin{cases} \rho^{\{\kappa_0\}} & \text{for } \kappa_0 \notin \mathbb{N}, \\ \rho \log(2 + \frac{1}{\rho}) & \text{for } \kappa_0 \in \mathbb{N}. \end{cases}$$

For every $|\beta| < \kappa_0$ we have

$$(31) \quad |D^\beta g(x)| \leq c|v - x|^{\alpha-d-|\beta|}, \quad x \neq v, |x| < 3/4.$$

Proof. Since $\text{supp}(\omega_B^v) \subset B^c$ the Proposition follows from (16), Theorem 1, (26), (27) and Lemma 4. $\square$

LEMMA 5 For every $0 < a < \gamma$ there exists a constant $c = c(a)$ such that

$$(32) \quad \int_{B(x, \rho)} |z - x|^{-a} \nu(dz) \leq c \rho^{\gamma-a} |x|^{-\alpha-\gamma}, \quad x \in \mathbb{R}^d \setminus \{0\}, 0 < \rho < \frac{|x|}{2}.$$

Proof. Let $x \in \mathbb{R}^d \setminus \{0\}$, $0 < \rho < |x|/2$. Then

$$\begin{aligned} \int_{B(x, \rho)} |z - x|^{-a} \nu(dz) &= \sum_{n=0}^{\infty} \int_{B(x, 2^{-n}\rho) \setminus B(x, 2^{-n-1}\rho)} |z - x|^{-a} \nu(dz) \\ &\leq \sum_{n=0}^{\infty} (2^{-n-1}\rho)^{-a} \nu(B(x, 2^{-n}\rho)) \\ &\leq \sum_{n=0}^{\infty} (2^{-n-1}\rho)^{-a} |x|^{-\alpha} \nu(B(x/|x|, 2^{-n}\rho/|x|)) \\ &\leq \sum_{n=0}^{\infty} (2^{-n-1}\rho)^{-a} c |x|^{-\alpha-\gamma} (2^{-n}\rho)^\gamma \\ &= \frac{2^a}{1 - 2^{a-\gamma}} c \rho^{\gamma-a} |x|^{-\alpha-\gamma}. \end{aligned}$$

$\square$

We define

$$s(x) = E^x \tau_{B(0,1)} = \int_{B(0,1)} G(x, v) dv.$$

The following result of M. Lewandowski is consequence of symmetry and nondegeneracy of the spectral measure μ (for the proof see [12, Lemma 10]).

LEMMA 6 *There exists c such that*

$$s(x) \leq c(1 - |x|^2)^{\alpha/2}, \quad |x| < 1.$$

LEMMA 7 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ then*

$$(33) \quad G_B(x, v) \leq c(1 - |v|)^{\alpha/2}, \quad |x| < 1/2, |v| > 3/4.$$

Proof. We refer to the proof of Lemma 16 in [13]. The only modification we need to make is Lemma 5 above with $a = d - \alpha$ to estimate $U_r g_n(v)$ in [13].

□

5 Poisson kernel

LEMMA 8 *There exists a constant c such that for every $r > 0$ we have*

$$(34) \quad \nu(B(z, r)) \leq cr^\gamma |z|^{-\gamma} (|z| - r)^{-\alpha}, \quad |z| > r.$$

Proof. For $|z| > 2r$ we have

$$\nu(B(z, r)) = |z|^{-\alpha} \nu(B(\frac{z}{|z|}, \frac{r}{|z|})) \leq cr^\gamma |z|^{-\alpha-\gamma},$$

whereas for $r < |z| \leq 2r$

$$\nu(B(z, r)) \leq \nu(B(0, |z| - r)^c) = c(|z| - r)^{-\alpha}.$$

□

LEMMA 9 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ then*

$$(35) \quad P_B(x, z) \leq c|z|^{-\gamma} (|z|^2 - 1)^{-\alpha/2}, \quad |x| < 1/2, |z| > 1.$$

Proof. By (19) and (17) we have

$$P_B(x, z) = \int_{B(z, 1)} G_B(x, z - v) \nu(dv) \leq \int_{B(z-x, 3/2)} c|v - (z - x)|^{\alpha-d} \nu(dv).$$

For $|z| > 3$ the estimate (35) follows from Lemma 5 with $a = d - \alpha$, and Lemma 8. For $1 < |z| < 3$ we have by (33), Lemma 5 with $a = d - \alpha$, and Lemma 8,

$$\begin{aligned}
P_B(x, z) &= \int_{B(z, 1)} G_B(x, z - v) \nu(dv) \\
&\leq \int_{B(z, 1) \setminus B(z, 3/4)} c(1 - |z - v|)^{\alpha/2} \nu(dv) + \int_{B(z, 3/4)} c|v - (z - x)|^{\alpha-d} \nu(dv) \\
&\leq \int_{B(0, |z|-1)^c} |v|^{\alpha/2} \nu(dv) + c \\
&\leq c(|z| - 1)^{-\alpha/2}.
\end{aligned}$$

□

The main result of this section is the following lemma.

LEMMA 10 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ and $\kappa_1 \neq 1$ then for every $|z| > 1$ the function $k_z(x) = P_B(x, z)$ belongs to the Hölder space $C^{\kappa_1}(B(0, 1/2))$. If $\kappa_1 = 1$ then $k_z(x) \in C^{1-}(B(0, 1/2))$. Moreover there is a constant c such that for every $|\beta| = [\kappa_1]^-$ we have*

$$\begin{aligned}
(36) \quad |D^\beta k_z(y) - D^\beta k_z(x)| &\leq cf(|y - x|)|z|^{-\gamma}(|z| - 1)^{-\alpha}, \\
&\quad x, y \in B(0, 1/2), |z| > 1,
\end{aligned}$$

where

$$f(\rho) = \begin{cases} \rho^{\{\kappa_1\}} & \text{for } \kappa_1 \notin \mathbb{N}, \\ \rho \log(\frac{2}{\rho}) & \text{for } \kappa_1 \in \mathbb{N}. \end{cases}$$

If $\kappa_1 > 1$ then for every $i \in \{1, \dots, d\}$ we have

$$(37) \quad |D_i k_z(x)| \leq c|z|^{-\gamma}(|z| - 1)^{-\alpha}, \quad |x| < 1/2, |z| > 1.$$

Proof. We assume first that $\kappa_1 = \gamma - (d - \alpha) > 1$. For every fixed $x \in B(0, 1/2)$, $i \in \{1, \dots, d\}$ and $|z| > 1$ we denote

$$f_h(v) = \frac{G_B(x + he_i, z - v) - G_B(x, z - v)}{h}, \quad h \in \mathbb{R}, v \in B,$$

where $\{e_1, \dots, e_d\}$ is the standard orthonormal basis in $\mathbb{R}^d$. We will prove that the family of functions $\{f_h(v), |h| < 1/16\}$ is uniformly integrable with respect

to the measure $\mathbf{1}_{B(z,1)}\nu(dv)$. Thus we need to estimate the integral

$$\begin{aligned}\int_{B(z,1)} |f_h(v)|^{1+\varepsilon} \nu(dv) &= \int_{B(z-x,2|h|)} + \int_{B(z-x,1/8) \setminus B(z-x,2|h|)} + \int_{B(z,1) \setminus B(z-x,1/8)} \\ &= A + B + C,\end{aligned}$$

where $0 < \varepsilon < \frac{\gamma}{d-\alpha+1} - 1$. By Lemma 5 with $a = (d-\alpha)(1+\varepsilon)$ and (17) we have

$$\begin{aligned}A &\leq \frac{2^{1+\varepsilon}}{|h|^{1+\varepsilon}} \left[\int_{B(z-x-he_i,3|h|)} (G_B(x+he_i, z-v))^{1+\varepsilon} \nu(dv) \right. \\ &\quad \left. + \int_{B(z-x,3|h|)} (G_B(x, z-v))^{1+\varepsilon} \nu(dv) \right] \\ &\leq \frac{2^{1+\varepsilon}}{|h|^{1+\varepsilon}} \left[\int_{B(z-x-he_i,3|h|)} c|z-x-he_i-v|^{(\alpha-d)(1+\varepsilon)} \nu(dv) \right. \\ &\quad \left. + \int_{B(z-x,3|h|)} c|z-x-v|^{(\alpha-d)(1+\varepsilon)} \nu(dv) \right] \\ &\leq \frac{c}{|h|^{1+\varepsilon}} |h|^{\gamma+(\alpha-d)(1+\varepsilon)} |z|^{-\alpha-\gamma} \\ &\leq c|z|^{-\alpha-\gamma}.\end{aligned}$$

Let $L = \lceil -\log_2(8|h|) \rceil$. By (31) with $|\beta| = 1$ and Lagrange's theorem we obtain

$$\begin{aligned}B &\leq \sum_{n=1}^L \int_{B(z-x,2^{n+1}|h|) \setminus B(z-x,2^n|h|)} |f_h(v)|^{1+\varepsilon} \nu(dv) \\ &\leq \sum_{n=1}^L c(2^{n-1}|h|)^{(\alpha-d-1)(1+\varepsilon)} \nu(B(z-x, 2^{n+1}|h|)) \\ &\leq \sum_{n=1}^L c(2^{n-1}|h|)^{(\alpha-d-1)(1+\varepsilon)} |z-x|^{-\alpha} \nu\left(B\left(\frac{z-x}{|z-x|}, \frac{2^{n+1}|h|}{|z-x|}\right)\right) \\ &\leq \sum_{n=1}^L c(2^{n-1}|h|)^{(\alpha-d-1)(1+\varepsilon)} (2^{n+1}|h|)^\gamma |z-x|^{-\alpha-\gamma} \\ &= c|h|^{(\alpha-d-1)(1+\varepsilon)+\gamma} |z-x|^{-\alpha-\gamma} \sum_{n=1}^L (2^{(\alpha-d-1)(1+\varepsilon)+\gamma})^n \\ &\leq c|z|^{-\alpha-\gamma}.\end{aligned}$$

Also by (31) we get

$$\begin{aligned} C &= \int_{B(z,1) \setminus B(z-x,1/8)} |f_h(v)|^{1+\varepsilon} \nu(dv) \\ &\leq \int_{B(z,1) \setminus B(z-x,1/8)} c\nu(dv) \leq c\nu(B(z,1)). \end{aligned}$$

We obtained

$$\int_{B(z,1)} |f_h(v)|^{1+\varepsilon} \nu(dv) = A + B + C \leq c(|z|^{-\alpha-\gamma} + \nu(B(z,1))),$$

which yields the uniform integrability of the family of functions $\{f_h(v)\}$. This, together with Proposition 1 and (19), yields

$$(38) \quad D_{x_i} k_z(x) = \int_{B(z,1)} D_{x_i} G_B(x, z-v) \nu(dv), \quad |x| < 1/2, |z| > 1,$$

and by (31), Lemma 5 with $a = d - \alpha + 1$ and Lemma 8 we get also (37) (compare the proof of Lemma 9).

Let now $\kappa_1 > 0$, $\kappa_1 \neq 1$ and $|\beta| = [\kappa_1] = [\kappa_1]^-$. We note that $\kappa_1 < 2$ so that $|\beta| \in \{0, 1\}$. Let $|z| > 1$, $x, y \in B(0, 1/2)$ and $|y - x| < 1/16$. We have

$$\begin{aligned} |D^\beta k_z(y) - D^\beta k_z(x)| &\leq \int_{B(z,1)} |D_x^\beta G_B(y, z-v) - D_x^\beta G_B(x, z-v)| \nu(dv) \\ &= \int_{B(z-x, 2|y-x|)} + \int_{B(z-x, 1/8) \setminus B(z-x, 2|y-x|)} \\ &\quad + \int_{B(z,1) \setminus B(z-x, 1/8)} \\ &= I + II + III. \end{aligned}$$

By (31) and Lemma 5 with $a = d - \alpha + |\beta|$ we have

$$\begin{aligned} I &\leq \int_{B(z-y, 3|y-x|)} |D_x^\beta G_B(y, z-v)| \nu(dz) \\ &\quad + \int_{B(z-x, 3|y-x|)} |D_x^\beta G_B(x, z-v)| \nu(dv) \\ &\leq \int_{B(z-y, 3|y-x|)} c|z - y - v|^{\alpha-d-|\beta|} \nu(dv) \end{aligned}$$

$$\begin{aligned}
& + \int_{B(z-x, 3|y-x|)} c|z-x-v|^{\alpha-d-|\beta|} \nu(dv) \\
& \leq c|y-x|^{\{\kappa_1\}} |z-y|^{-\alpha-\gamma} + c|y-x|^{\{\kappa_1\}} |z-x|^{-\alpha-\gamma} \\
& \leq c|y-x|^{\{\kappa_1\}} |z|^{-\alpha-\gamma}.
\end{aligned}$$

In order to estimate *II* and *III* we assume first $\kappa_0 \notin \mathbb{N}$ or $\kappa_0 - \kappa_1 = \alpha > 1$. Let $\delta_0 = \min(\kappa_0, |\beta| + 1)$. Then we have $[\delta_0]^- = |\beta| = [\kappa_1]$, because $\kappa_0 > \kappa_1 > [\kappa_1]$. We use (30) in case $\delta_0 = \kappa_0$, and (31) and Lagrange's theorem in case $\delta_0 = |\beta| + 1$, to obtain for every $r \in (0, 1/2)$ and $w \in B$

$$(39) \quad |D_x^\beta G_B(y, w) - D_x^\beta G_B(x, w)| \leq cr^{\alpha-d-\delta_0} |y-x|^{\delta_0-[\delta_0]^-},$$

provided $x, y \in B(0, 1/2) \cap B(w, r)^c$. Hence for $L = \lceil -\log_2(8|y-x|) \rceil$ we get

$$\begin{aligned}
II & \leq \sum_{n=1}^L \int_{B(z-x, 2^{n+1}|y-x|) \setminus B(z-x, 2^n|y-x|)} |D_x^\beta G_B(y, z-v) - D_x^\beta G_B(x, z-v)| \nu(dv) \\
& \leq \sum_{n=1}^L c(2^{n-1}|y-x|)^{\alpha-d-\delta_0} |y-x|^{\delta_0-[\delta_0]^-} \nu(B(z-x, 2^{n+1}|y-x|)) \\
& \leq \sum_{n=1}^L c(2^{n-1})^{\alpha-d-\delta_0} |y-x|^{\alpha-d-[\delta_0]^-} |z-x|^{-\alpha} \nu\left(B\left(\frac{z-x}{|z-x|}, \frac{2^{n+1}|y-x|}{|z-x|}\right)\right) \\
& \leq \sum_{n=1}^L c(2^{n-1})^{\alpha-d-\delta_0} |y-x|^{\alpha-d-[\delta_0]^-} (2^{n+1}|y-x|)^\gamma |z-x|^{-\alpha-\gamma} \\
& \leq c|y-x|^{\alpha+\gamma-d-[\delta_0]^-} |z-x|^{-\alpha-\gamma} \sum_{n=1}^{\infty} (2^{\alpha-d-\delta_0+\gamma})^n \\
& \leq c|y-x|^{\kappa_1-[\delta_0]^-} |z|^{-\alpha-\gamma} \\
& = c|y-x|^{\{\kappa_1\}} |z|^{-\alpha-\gamma},
\end{aligned}$$

and

$$\begin{aligned}
III & = \int_{B(z, 1) \setminus B(z-x, 1/8)} |D_x^\beta G_B(y, z-v) - D_x^\beta G_B(x, z-v)| \nu(dv) \\
& \leq \int_{B(z, 1) \setminus B(z-x, 1/8)} c(1/16)^{\alpha-d-\delta_0} |y-x|^{\delta_0-[\delta_0]^-} \nu(dv) \\
& \leq c|y-x|^{\delta_0-[\delta_0]^-} \nu(B(z, 1)) \\
& \leq c|y-x|^{\delta_0-[\delta_0]^-} |z|^{-\gamma} (|z|-1)^{-\alpha}.
\end{aligned}$$

This, together with Lemma 9 and (37), yield (36) for $\kappa_0 \notin \mathbb{N}$ or $\kappa_0 - \kappa_1 > 1$.
 If $\kappa_0 \in \mathbb{N}$ and $\kappa_0 - \kappa_1 < 1$ then by (30) we obtain

$$\begin{aligned}
 II &\leq \sum_{n=1}^L \int_{B(z-x, 2^{n+1}|y-x|) \setminus B(z-x, 2^n|y-x|)} |D_x^\beta G_B(y, z-v) - D_x^\beta G_B(x, z-v)| \nu(dv) \\
 &\leq \sum_{n=1}^L c(2^{n-1}|y-x|)^{\alpha-d-|\beta|} 2^{-n+1} \log(2^{n+1}) \nu(B(z-x, 2^{n+1}|y-x|)) \\
 &\leq \sum_{n=1}^L cn(2^{n-1})^{\alpha-d-|\beta|-1} |y-x|^{\alpha-d-|\beta|} |z-x|^{-\alpha} \nu(B(\frac{z-x}{|z-x|}, \frac{2^{n+1}|y-x|}{|z-x|})) \\
 &\leq \sum_{n=1}^L cn(2^{n-1})^{\alpha-d-|\beta|-1} |y-x|^{\alpha-d-|\beta|} (2^{n+1}|y-x|)^\gamma |z-x|^{-\alpha-\gamma} \\
 &\leq c|y-x|^{\kappa_1-|\beta|} |z-x|^{-\alpha-\gamma} \sum_{n=1}^{\infty} n(2^{-\alpha})^n \\
 &\leq c|y-x|^{\{\kappa_1\}} |z|^{-\alpha-\gamma},
 \end{aligned}$$

and

$$\begin{aligned}
 III &= \int_{B(z,1) \setminus B(z-x,1/8)} |D_x^\beta G_B(y, z-v) - D_x^\beta G_B(x, z-v)| \nu(dv) \\
 &\leq \int_{B(z,1) \setminus B(z-x,1/8)} c(1/16)^{\alpha-d-\kappa_0} \log(2 + \frac{1}{16|y-x|}) |y-x| \nu(dv) \\
 &\leq c \log(2 + \frac{1}{16|y-x|}) |y-x| \nu(B(z,1)) \\
 &\leq c|y-x|^{\{\kappa_1\}} |z|^{-\gamma} (|z|-1)^{-\alpha},
 \end{aligned}$$

and we get (36) also in this case.

If $\kappa_1 = 1$ and $\beta = 0$ then by Lemma 5 with $a = d - \alpha$, (31) and Lagrange's theorem we get

$$\begin{aligned}
 I &\leq \int_{B(z-y, 3|y-x|)} |G_B(y, z-v)| \nu(dz) \\
 &\quad + \int_{B(z-x, 3|y-x|)} |G_B(x, z-v)| \nu(dv) \\
 &\leq c|y-x||z|^{-\alpha-\gamma},
 \end{aligned}$$

$$\begin{aligned}
II &\leq c|y-x|^{\alpha+\gamma-d}|z-x|^{-\alpha-\gamma} \sum_{n=1}^L (2^{\alpha-d-1+\gamma})^n \\
&= c|y-x||z-x|^{-\alpha-\gamma} L \\
&\leq c|y-x||z-x|^{-\alpha-\gamma} \log(|y-x|^{-1}),
\end{aligned}$$

and

$$\begin{aligned}
III &= \int_{B(z,1) \setminus B(z-x,1/8)} |G_B(y, z-v) - G_B(x, z-v)| \nu(dv) \\
&\leq c|y-x||z|^{-\gamma}(|z|-1)^{-\alpha},
\end{aligned}$$

which yields (36) also in this case. By (35), (37) and (36) we get $k_z \in C^{\kappa_1}(B(0, 1/2))$ for $\kappa_1 \neq 1$ and $k_z \in C^p(B(0, 1/2))$ for $\kappa_1 = 1$ and $p \in (0, 1)$.

□

6 Harmonic functions

The Hölder continuity of harmonic functions of stable processes under assumption that ν is absolutely continuous was first proved in [2]. Here we do not generally assume that ν is absolutely continuous (or $\gamma = d$). However, if so is the case, then we obtain $\kappa_1 = \alpha$ and we get the α -Hölder continuity for $\alpha < 1$ and the existence of the first derivative of bounded harmonic functions for $\alpha > 1$.

The following lemma contains continuity estimates for bounded harmonic functions. These inequalities yield Hölder continuity.

LEMMA 11 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ then there exists a constant c such that for every function u bounded on $\mathbb{R}^d$ and harmonic in B we have*

$$|u(y) - u(x)| \leq c \|u\|_{\infty} f(|y - x|), \quad x, y \in B(0, 1/2),$$

where

$$f(\rho) = \begin{cases} \rho^{\kappa_1} & \text{for } \alpha < 1, \\ \rho^{\kappa_1} \log(2/\rho) & \text{for } \alpha = 1 \text{ and } \gamma < d, \\ \rho \log^2(2/\rho) & \text{for } \alpha = 1 \text{ and } \gamma = d, \\ \rho^{\frac{2-\alpha}{\alpha} \min(\kappa_1, 1)} & \text{for } \alpha > 1 \text{ and } \kappa_1 \neq 1, \\ [\rho \log(2/\rho)]^{\frac{2-\alpha}{\alpha}} & \text{for } \alpha > 1 \text{ and } \kappa_1 = 1. \end{cases}$$

Proof. Let $\delta_0 = \min(\kappa_1, 1)$. If $\kappa_1 > 0$ and $\kappa_1 \neq 1$ then by Lemma 10, 9 and Lagrange's theorem for $\alpha > 1$ we obtain

$$\begin{aligned}
|u(y) - u(x)| &\leq \int_{B^c} |u(z)| |P_B(y, z) - P_B(x, z)| dz \\
&\leq \|u\|_\infty \int_{B(0,2)^c} c|y-x|^{\delta_0} |z|^{-\gamma-\alpha} dz \\
&\quad + \|u\|_\infty \int_{B(0,2) \setminus B(0,1+|y-x|^{2\delta_0/\alpha})} c|y-x|^{\delta_0} (|z|-1)^{-\alpha} dz \\
&\quad + \|u\|_\infty \int_{B(0,1+|y-x|^{2\delta_0/\alpha}) \setminus B(0,1)} c(|z|-1)^{-\alpha/2} dz \\
&= c\|u\|_\infty (|y-x|^{\delta_0} + |y-x|^{\delta_0 \frac{2-\alpha}{\alpha}}) \\
&\leq c\|u\|_\infty |y-x|^{\delta_0 \frac{2-\alpha}{\alpha}}.
\end{aligned}$$

For $\alpha < 1$ by Lemma 10 we have

$$\begin{aligned}
|u(y) - u(x)| &\leq \int_{B^c} |u(z)| |P_B(y, z) - P_B(x, z)| dz \\
&\leq \|u\|_\infty \int_{B^c} c|y-x|^{\delta_0} |z|^{-\gamma} (|z|-1)^{-\alpha} dz \\
&= c\|u\|_\infty |y-x|^{\delta_0}.
\end{aligned}$$

Similarly for $\alpha = 1$ we get

$$|u(y) - u(x)| \leq c\|u\|_\infty |y-x|^{\delta_0} \log\left(\frac{2}{|y-x|}\right).$$

Let $r = \lceil \log(2|y-x|^{-1}) \rceil |y-x|^{2/\alpha}$. If $\kappa_1 = 1$ then for $\alpha > 1$ by Lemma 10 we obtain

$$\begin{aligned}
|u(y) - u(x)| &\leq \int_{B^c} |u(z)| |P_B(y, z) - P_B(x, z)| dz \\
&\leq \|u\|_\infty \int_{B(0,2)^c} c \log(2|y-x|^{-1}) |y-x| |z|^{-\gamma-\alpha} dz \\
&\quad + \|u\|_\infty \int_{B(0,2) \setminus B(0,1+r))} c \log(2|y-x|^{-1}) |y-x| (|z|-1)^{-\alpha} dz \\
&\quad + \|u\|_\infty \int_{B(0,1+r) \setminus B(0,1)} c(|z|-1)^{-\alpha/2} dz
\end{aligned}$$

$$\begin{aligned}
&\leq c\|u\|_\infty \left[\log(2|y-x|^{-1})|y-x| + [\log(2|y-x|^{-1})|y-x|]^{\frac{2-\alpha}{\alpha}} \right] \\
&\leq c\|u\|_\infty [\log(2|y-x|^{-1})|y-x|]^{\frac{2-\alpha}{\alpha}},
\end{aligned}$$

and for $\alpha = 1$, $\gamma = d$ we have

$$\begin{aligned}
|u(y) - u(x)| &\leq \int_{B^c} |u(z)| |P_B(y, z) - P_B(x, z)| dz \\
&\leq \|u\|_\infty \int_{B(0,2)^c} c \log(2|y-x|^{-1})|y-x| |z|^{-d-1} dz \\
&\quad + \|u\|_\infty \int_{B(0,2) \setminus B(0,1+r)} c \log(2|y-x|^{-1})|y-x| (|z| - 1)^{-1} dz \\
&\quad + \|u\|_\infty \int_{B(0,1+r) \setminus B(0,1)} c (|z| - 1)^{-1/2} dz \\
&\leq c\|u\|_\infty \log(2|y-x|^{-1})|y-x| \log [(\log(2|y-x|^{-1})|y-x|)^{-1}], \\
&\leq c\|u\|_\infty \log^2(2|y-x|^{-1})|y-x|.
\end{aligned}$$

□

Note that translation invariance and the scaling property of the process yields that if the function $y \rightarrow u(y)$ is harmonic in $B(z, r)$ then the function $y \rightarrow u(ry + z)$ is harmonic in B . Therefore we have

$$(40) \quad |u(y) - u(x)| \leq c\|u\|_\infty f(|y-x|/r), \quad x, y \in B(z, r/2),$$

for every $z \in \mathbb{R}^d$, $r > 0$ and bounded function u harmonic in $B(z, r)$.

LEMMA 12 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ then there exists a constant c such that for every bounded open set D we have*

$$(41) \quad P_D(x, z) \leq c \operatorname{dist}(x, D^c)^{\alpha-d} \operatorname{dist}(z, D)^{-\alpha}, \quad x \in D, z \in D^c.$$

Proof. Let $x \in D$ and $z \in D^c$. By (17), (19) and Lemma 5 we have

$$\begin{aligned}
P_D(x, z) &= \int_{z-D} G_D(x, z-v) \nu(dv) \\
&\leq c \int_{z-D} |z-v-x|^{\alpha-d} \nu(dv) \\
&\leq c \int_{B(z-x, \operatorname{dist}(x, D^c)/2)} |z-v-x|^{\alpha-d} \nu(dv)
\end{aligned}$$

$$\begin{aligned}
& +c \int_{z-D} (\text{dist}(x, D^c)/2)^{\alpha-d} \nu(dv) \\
& \leq c[\text{dist}(x, D^c)]^{\gamma+\alpha-d} |z-x|^{-\alpha-\gamma} \\
& \quad +c(\text{dist}(x, D^c))^{\alpha-d} \nu(B(0, \text{dist}(z, D))^c) \\
& \leq c \text{dist}(x, D^c)^{\alpha-d} \text{dist}(z, D)^{-\alpha}.
\end{aligned}$$

□

LEMMA 13 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ and $\alpha > 1$ then for every $i \in \{1, \dots, d\}$ the function $D_i V(x)$ is harmonic in $\mathbb{R}^d \setminus \{0\}$.*

Proof. Let $R > r > 0$, $h \in (-r/16, r/16)$ and $x \in A = B(0, R) \setminus B(0, r)$. Since the potential kernel $V(x)$ is harmonic in $\mathbb{R}^d \setminus \{0\}$ we have

$$V(x + he_i) = E^{x+he_i} V(X_{\tau_{(A+he_i)}}) = E^x V(X_{\tau_A} + he_i),$$

and

$$\begin{aligned}
\frac{1}{h}(V(x + he_i) - V(x)) &= \frac{1}{h} E^x [V(X_{\tau_A} + he_i) - V(X_{\tau_A})] \\
&= \frac{1}{h} \int_{A^c} [V(z + he_i) - V(z)] \omega_A^x(dz).
\end{aligned}$$

Let

$$f_h(z) = \frac{V(z + he_i) - V(z)}{h}.$$

We will prove that the family of functions $\{f_h(z), |h| < r/16\}$ is uniformly integrable with respect to the measure ω_A^x . To this end we will estimate the integral

$$\begin{aligned}
\int_{A^c} |f_h(z)|^{1+\varepsilon} \omega_A^x(dz) &= \int_{B(0, 2|h|)} + \int_{B(0, r/8) \setminus B(0, 2|h|)} + \int_{A^c \setminus B(0, r/8)} \\
&= I + II + III,
\end{aligned}$$

where $0 < \varepsilon < \frac{d}{d-\alpha+1} - 1$. By Lemma 12 we have

$$\begin{aligned}
I &\leq \frac{2^{1+\varepsilon}}{|h|^{1+\varepsilon}} \left[\int_{B(-he_i, 3|h|)} (V(z + he_i))^{1+\varepsilon} \omega_A^x(dz) + \int_{B(0, 3|h|)} (V(z))^{1+\varepsilon} \omega_A^x(dz) \right] \\
&\leq \frac{2^{1+\varepsilon}}{|h|^{1+\varepsilon}} \left[\int_{B(-he_i, 3|h|)} c|z + he_i|^{(\alpha-d)(1+\varepsilon)} P_A(x, z) dz \right]
\end{aligned}$$

$$\begin{aligned}
& + \int_{B(0,3|h|)} c|z|^{(\alpha-d)(1+\varepsilon)} P_A(x, z) dz \\
& \leq \frac{c}{|h|^{1+\varepsilon}} |h|^{d+(\alpha-d)(1+\varepsilon)} \text{dist}(x, A^c)^{\alpha-d} (3r/4)^{-\alpha} \\
& \leq c \text{dist}(x, A^c)^{\alpha-d} (3r/4)^{-\alpha}.
\end{aligned}$$

By (1) and Lagranger's theorem for $L = \lceil \log_2(\frac{r}{8|h|}) \rceil$ we obtain

$$\begin{aligned}
II & \leq \sum_{n=1}^L \int_{B(0,2^{n+1}|h|) \setminus B(0,2^n|h|)} |f_h(z)|^{1+\varepsilon} \omega_A^x(dz) \\
& \leq \sum_{n=1}^L c(2^{n-1}|h|)^{(\alpha-d-1)(1+\varepsilon)} \omega_A^x(B(0, 2^{n+1}|h|)) \\
& \leq \sum_{n=1}^L c(2^{n-1}|h|)^{(\alpha-d-1)(1+\varepsilon)} \text{dist}(x, A^c)^{\alpha-d} (3r/4)^{-\alpha} (2^{n+1}|h|)^d \\
& = c|h|^{(\alpha-d-1)(1+\varepsilon)+d} \text{dist}(x, A^c)^{\alpha-d} (3r/4)^{-\alpha} \sum_{n=1}^L (2^{(\alpha-d-1)(1+\varepsilon)+d})^n \\
& \leq c \text{dist}(x, A^c)^{\alpha-d} r^{(\alpha-d-1)(1+\varepsilon)+d-\alpha},
\end{aligned}$$

$$\begin{aligned}
III & = \int_{A^c \setminus B(0,r/8)} |f_h(z)|^{1+\varepsilon} \omega_A^x(dz) \\
& \leq \int_{A^c \setminus B(0,r/8)} c r^{(\alpha-d-1)(1+\varepsilon)} \omega_A^x(dz) \\
& \leq c r^{(\alpha-d-1)(1+\varepsilon)} \omega_A^x(A^c) = c r^{(\alpha-d-1)(1+\varepsilon)}.
\end{aligned}$$

We obtained

$$\int_{A^c} |f_h(z)|^{1+\varepsilon} \omega_A^x(dz) = I + II + III \leq c(r, R, x)$$

which clearly yields the uniform integrability of the family of functions $\{f_h(v)\}$.

We obtain

$$\begin{aligned}
D_i V(x) & = \lim_{h \rightarrow 0} \frac{V(x + h e_i) - V(x)}{h} \\
& = \lim_{h \rightarrow 0} \int_{A^c} \frac{V(z + h e_i) - V(z)}{h} \omega_A^x(dz)
\end{aligned}$$

$$\begin{aligned}
&= \int_{A^c} D_i V(z) \omega_A^x(dz) \\
&= E^x D_i V(X_{\tau_A}).
\end{aligned}$$

This yields regular harmonicity of $D_i V$ in $A = B(0, R) \setminus B(0, r)$ for every $R > r > 0$ and harmonicity in $\mathbb{R}^d \setminus \{0\}$. $\square$

COROLLARY 14 *If $\kappa_1 = \gamma - (d - \alpha) > 0$ and $\alpha > 1$ then for every $i \in \{1, \dots, d\}$ and $x \in B$ the function $g(v) = D_{x_i} G_B(x, v)$ is harmonic in $B \setminus \{x\}$ and continuous in $\mathbb{R}^d \setminus \{x\}$.*

Proof. By symmetry of the Green function, (16) and Lemma 4 we have

$$(42) \quad D_{x_i} G_B(x, v) = -D_i V(v - x) - E^v D_i V(x - X_{\tau_B})$$

which is harmonic in $B \setminus \{x\}$ by Lemma 13. The function $v \rightarrow D_i V(v - x)$ is continuous in $\mathbb{R}^d \setminus \{x\}$ by Theorem 1. The function

$$h(v) = E^v D_i V(x - X_{\tau_B}), \quad v \in B,$$

is bounded and harmonic in $B(v_0, 1 - |v_0|)$ for every $v_0 \in B$ hence from (40) we obtain the continuity of $h(v)$ at v_0 and the continuity of $v \rightarrow D_{x_i} G_B(x, v)$ in B . $D_{x_i} G_B(x, v) = 0$ for $v \in B^c$ and we have also $\lim_{v \rightarrow w} D_{x_i} G_B(x, v) = 0$ for every $x \in B(0, 1)$ and every point $w \in \mathbb{S}$ because the measures ω_B^v weakly converge to δ_w (see, e.g., [16, Theorem 1.23]). $\square$

We will employ the operator

$$\mathcal{U}_r \phi(x) = \frac{E^x \phi(X_{\tau_{B(x, r)}}) - \phi(x)}{E^x \tau_{B(x, r)}},$$

whenever the expression is well defined for given ϕ , $r > 0$ and x . Clearly, if h is harmonic in D , $x \in D$, and $r < \text{dist}(x, D^c)$, then $\mathcal{U}_r h(x) = 0$. We note that

$$\mathcal{U} \phi(x) = \lim_{r \downarrow 0} \mathcal{U}_r \phi(x)$$

is the Dynkin characteristic operator, which was used in [13] in a similar way.

We record the following observation (maximum principle).

LEMMA 15 *If there is $r > 0$ such that $\mathcal{U}_r h(x) > 0$ then $h(x) < \sup_{y \in \mathbb{R}^d} h(y)$.*

The proof of the following Lemma is based on ideas of [13, Lemma 16].

LEMMA 16 *If $\kappa_1 = \gamma - (d - \alpha) > 1$ then there exists c such that*

$$(43) \quad |D_{x_i} G_B(x, v)| < c(1 - |v|)^{\alpha/2}, \quad |x| < 1/2, \quad 3/4 < |v| < 1.$$

Proof. By the strong Markov property we have

$$\begin{aligned} s(v) &= E^v \tau_B = E^v(\tau_A + \tau_{B(0,1)} \circ \theta_{\tau_A}) = E^v \tau_A + E^v E^{X_{\tau_A}} \tau_{B(0,1)} \\ &= E^v \tau_A + E^v s(X_{\tau_A}), \quad v \in \mathbb{R}^d, \quad A \subset B(0, 1), \end{aligned}$$

which yields $\mathcal{U}_r s(v) = -1$ for $v \in B(0, 1)$ and $r < 1 - |v|$.

For $n \in \mathbb{N}$ and $x \in B(0, 1/2)$ we let $g(v) = D_{x_i} G_B(x, v)$, $g_{1,n}(v) = \min(g(v), n)$ and $g_{2,n}(v) = \max(g(v), -n)$. For $v \in B(x, 1/8)^c$ we have that $|D_{x_i} G_B(x, v)| \leq c_1 |v - x|^{\alpha-d-1}$ hence $g_{1,n}(v) = g_{2,n}(v) = g(v)$ provided $n \geq c_1 8^{d-\alpha+1}$. It follows from Corollary 14 that $g(v)$ is harmonic on $B \setminus \{x\}$. Hence by scaling property, (18), Lemma 5 and 8 we obtain that for $v \in B \setminus B(0, 3/4)$ and $r < \min(1 - |v|, 1/16)$ it holds

$$\begin{aligned} \mathcal{U}_r g_{1,n}(v) &= \mathcal{U}_r(g_{1,n} - g)(v) \\ &= \frac{1}{E^0 \tau_B} \int_B G(0, w) \int (g_{1,n} - g)(v + rw + z) \nu(dz) dw \\ &\geq \frac{-c_2}{s(0)} \int_B G(0, w) \int_{B(x-v-rw, 1/8)} |x - v - rw - z|^{\alpha-d-1} \nu(dz) dw \\ &\geq -c_3, \end{aligned}$$

Similarly

$$\begin{aligned} \mathcal{U}_r g_{2,n}(v) &= \mathcal{U}_r(g_{2,n} - g)(v) \\ &\leq c_4, \end{aligned}$$

If $a > c_3$ and $a > c_4$ then

$$\mathcal{U}_r(as - g_{1,n})(v) = -a - \mathcal{U}_r g_{1,n}(v) \leq -a + c_3 < 0,$$

and

$$\mathcal{U}_r(as + g_{2,n})(v) = -a + \mathcal{U}_r g_{2,n}(v) \leq -a + c_4 < 0.$$

By scaling

$$(44) \quad \begin{aligned} s(v) &\geq E^v \tau_{B(v, 1-|v|)} = (1-|v|)^\alpha E^0 \tau_{B(0,1)} \\ &\geq 4^{-\alpha} E^0 \tau_B, \quad |v| < 3/4. \end{aligned}$$

Since $g_{1,n}(v) \leq n$ and $g_{2,n}(v) \geq -n$, we see that $as(v) - g_{1,n}(v) > 0$ and $as(v) + g_{2,n}(v) > 0$ for $v \in B(0, 3/4)$ provided $a > n/(4^{-\alpha} E^0 \tau_B)$.

Let $a_0 = \max[c_3, c_4, n/(4^{-\alpha} E^0 \tau_{B(0,1)})] + 1$ and $h_1(v) = a_0 s(v) - \overline{g_{1,n}(v)}$, $h_2(v) = a_0 s(v) + g_{2,n}(v)$. We have $h_1(v) \geq 0$ and $h_2(v) \geq 0$ for $v \in \overline{B(0, 3/4)}$, $h_1(v) = h_2(v) = 0$ for $v \in B(0, 1)^c$ and $\mathcal{U}_r h_1(v) < 0$, $\mathcal{U}_r h_2(v) < 0$ for $v \in B(0, 1) \setminus B(0, 3/4)$, $r < \min(1-|v|, 1/16)$. Lemma 15 and continuity of h_1 and h_2 in $\mathbb{R}^d \setminus \{x\}$ yields $h_1(v) \geq 0$ and $h_2(v) \geq 0$ in $B(0, 1)$. Since $g_{1,n} = g_{2,n} = g$ on $B(0, 3/4)^c$, the Lemma follows from Lemma 6. $\square$

Using (43) we obtain the following estimate. We omit the proof since it is analogous to the proof of Lemma 9

LEMMA 17 *If $\kappa_1 = \gamma - (d - \alpha) > 1$ then there exists a constant c such that for every $i \in \{1, \dots, d\}$ we have*

$$(45) \quad |D_{x_i} P_B(x, z)| \leq c |z|^{-\gamma} (|z|^2 - 1)^{-\alpha/2}, \quad |x| < 1/2, |z| > 1.$$

Now we can prove our main result.

Proof of Theorem 2. We assume first $\kappa_1 = \gamma - (d - \alpha) > 1$. By (45) we obtain

$$D_i u(x) = \int_{B^c} u(z) D_{x_i} P_B(x, z) dz, \quad x \in B(0, 1/2),$$

and by Lemma 10 and (45) we get

$$(46) \quad \begin{aligned} |D_i u(y) - D_i u(x)| &\leq \int_{B^c} |u(z)| |D_{x_i} P_B(y, z) - D_{x_i} P_B(x, z)| dz \\ &\leq \|u\|_\infty \int_{B(0, 2)^c} c |y - x|^{\{\kappa_1\}} |z|^{-\gamma-\alpha} dz \\ &\quad + \|u\|_\infty \int_{B(0, 2) \setminus B(0, 1+|y-x|^{2\{\kappa_1\}/\alpha})} c |y - x|^{\{\kappa_1\}} (|z| - 1)^{-\alpha} dz \\ &\quad + \|u\|_\infty \int_{B(0, 1+|y-x|^{2\{\kappa_1\}/\alpha}) \setminus B(0, 1)} c (|z| - 1)^{-\alpha/2} dz \\ &= c \|u\|_\infty (|y - x|^{\{\kappa_1\}} + |y - x|^{\{\kappa_1\} \frac{2-\alpha}{\alpha}}) \\ &\leq c \|u\|_\infty |y - x|^{\{\kappa_1\} \frac{2-\alpha}{\alpha}}, \quad x, y \in B(0, 1/2). \end{aligned}$$

This and Lemma 11 yield for every $\kappa_1 > 0$ some constant $\delta \in (0, 1)$ such that $u \in C^{[\kappa_1]^- + \delta}(B(0, 1/2))$. Let now $x_0 \in B$. The function $u(x)$ is harmonic in $B(x_0, 1 - |x_0|)$ therefore the function $h(x) = u((1 - |x_0|)x + x_0)$ is harmonic in B and for $|\beta| = [\kappa_1]^-$ we get

$$\begin{aligned} |D^\beta u(y) - D^\beta u(x_0)| &= (1 - |x_0|)^{-|\beta|} |(D^\beta h)\left(\frac{y - x_0}{1 - |x_0|}\right) - (D^\beta h)(0)| \\ &\leq c(1 - |x_0|)^{-|\beta| - \delta} |y - x_0|^\delta, \quad y \in B(x_0, \frac{1 - |x_0|}{2}), \end{aligned}$$

which clearly yields $u \in C_{loc}^{[\kappa_1]^- + \delta}(B)$ and (2). $\square$

Acknowledgments. The author is grateful to Prof. R. L. Schilling for useful discussions and his hospitality during the stay in Marburg and to Prof. K. Bogdan for discussions and suggestions on the paper.

References

- [1] R. F. Bass, M. Kassmann, *Hölder continuity of harmonic functions with respect to operators of variable orders*, Comm. Partial Differential Equations 30 (2005), 1249-1259.
- [2] R. F. Bass, D. A. Levin, *Harnack inequalities for jump processes*, Potential Anal. 17(4)(2002), 375-388.
- [3] J. Bertoin, *Lévy processes*, Cambridge University Press, Cambridge, 1996.
- [4] R. M. Blumenthal and R. K. Gettoor, *Markov Processes and Potential Theory*, Pure Appl. Math., Academic Press Inc., New York 1968.
- [5] K. Bogdan, T. Byczkowski, *Potential theory for the α -stable Schrödinger operator on bounded Lipschitz domains*, Studia Math. 133 (1999) no. 1, 53-92.
- [6] K. Bogdan, B. Dyda, *Relative Fatou theorem for harmonic functions of rotation invariant stable processes in smooth domains*, Stud. Math. 157, No. 1, 83-96 (2003).

- [7] K. Bogdan, T. Jakubowski, *Estimates of heat kernel of fractional Laplacian perturbed by gradient operators*, Comm. Math. Phys. 271 (1) 2007, 179–198.
- [8] K. Bogdan, T. Kulczycki, A. Nowak, *Gradient estimates for harmonic and q -harmonic functions of symmetric stable processes*, Ill. J. Math. 46, No. 2, 541–556 (2002).
- [9] K. Bogdan, T. Kulczycki, M. Kwaśnicki *Estimates and structure of α -harmonic functions*, Probab. Theory Relat. Fields 2007.
- [10] K. Bogdan, A. Stós, P. Sztonyk, *Potential theory for Lévy stable processes*, Bull. Polish. Acad. Sci. Math. 50(3) 2002, 361–372.
- [11] K. Bogdan, A. Stós, P. Sztonyk, *Harnack inequality for symmetric stable processes on fractals*, C. R. Math. Acad. Sci. Paris 335 (1) (2002), 59–63.
- [12] K. Bogdan, P. Sztonyk, *Harnack's inequality for stable Lévy processes*, Potential Analysis 22 (2) (2005), 133–150.
- [13] K. Bogdan, P. Sztonyk, *Estimates of potential kernel and Harnack's inequality for anisotropic fractional Laplacian*, to appear in Studia Math. (http://arxiv.org/PS_cache/math/pdf/0507/0507579.pdf).
- [14] K. L. Chung, *Lectures from Markov processes to Brownian motion*, Springer-Verlag, New York-Berlin, 1982.
- [15] K. L. Chung, *Doubly-Feller process with multiplicative functional*, Seminar on stochastic processes, 1985 (Gainesville, Fla., 1985), 63–78, Progr. Probab. Statist., 12, Birkhäuser Boston, Boston, MA, 1986.
- [16] K. L. Chung, Z. Zhao, *From Brownian motion to Schrödinger's equation*, Springer - Verlag, New York, 1995.
- [17] S.N. Ethier, Th. G. Kurtz, *Markov processes - characterization and convergence*, Wiley Series in Probability and Mathematical Statistics, John Wiley, New York - Chicester - Brisbane - Toronto - Singapore (1986).
- [18] E. De Giorgi, *Sulla differenziabilità e l'analiticità delle estremali degli integrali multipli regolari*, Mem. Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. (3), 3 (1957), 25–43.

- [19] M. Hardy, *Combinatorics of Partial Derivatives*, Electronic Journal of Combinatorics, 13 (2006).
- [20] S. Hiraba *Asymptotic estimates for densities of multi-dimensional stable distributions*, Tsukuba J. Math. 27 (2003), no. 2, 261–287.
- [21] B. Hu, H. -M. Yin, *The DeGiorgi-Nash-Moser type of estimate for parabolic Volterra integrodifferential equations*, Pacific J. Math., 178(2) (1997), 265–277.
- [22] R. Hussein, M. Kassmann, *Jump processes, $\mathcal{L}$ -harmonic functions, continuity estimates and the Feller property*, 2006, preprint.
- [23] N. Ikeda, S. Watanabe, *On some relations between the harmonic measure and the Lévy measure for a certain class of Markov processes*, J. Math. Kyoto Univ. 2-1 (1962), 79–95.
- [24] N. Jacob, *Pseudo differential operators and Markov processes. Vol. I. Fourier analysis and semigroups*, Imp. Coll. Press, London, 2001.
- [25] N. Jacob, *Pseudo-Differential Operators and Markov Processes, Vol. II : Generators and Their Potential Theory*, Imperial College Press, London, 2002.
- [26] T. Jakubowski, *The estimates for the Green function in Lipschitz domains for the symmetric stable processes*, Probab. Math. Statist. 22 (2002), 419–441.
- [27] M. Kassmann, *On regularity for Beurling-Deny type Dirichlet forms*, Potential Analysis 19 (2003), 69–87.
- [28] T. Kulczycki, *Properties of Green function of symmetric stable process*, Probab. Math. Statist. 17(2) (1997), 339–364.
- [29] M. Lewandowski *Point regularity of p -stable density in $\mathcal{R}^d$ and Fisher information*, Probab. Math. Stat. 19, No.2, 375–388 (1999).
- [30] K. Michalik, *Sharp estimates of the Green function, the Poisson kernel and the Martin kernel of cones for symmetric stable processes*, Hiroshima Math. J. 36, No. 1 (2006), 1–21.

- [31] K. Michalik, M. Ryznar, *Relative Fatou theorem for α -harmonic functions in Lipschitz domains*, Illinois J. Math. 48 (2004), 977-998.
- [32] R. Mikulevičius, H. Pragarauskas, *On Hölder continuity of solutions of certain integro-differential equations*, Ann. Acad. Sci. Fenn. Ser. A I Math., 13(2) (1988), 231-238.
- [33] P. W. Millar, *First passage distributions of processes with independent increments*, Ann. Probab., 3 (1975) no. 2, 215-233.
- [34] J. Moser, *On Harnack's theorem for elliptic differential equations*, Comm. Pure Appl. Math., 14 (1961), 577-591.
- [35] J. Nash, *Continuity of solutions of parabolic and elliptic equations*, Amer. J. Math., 80 (1958), 931-954.
- [36] J. Picard, *Density in small time at accessible points for jump processes*, Stochastic Process. Appl. 67 (1997), no. 2, 251-279.
- [37] W. Rudin, *Functional Analysis*, 2nd ed., International Series in Pure and Applied Mathematics. New York, NY: McGraw-Hill. xviii, 424 p. (1991).
- [38] K.-I. Sato, *Lévy Processes and Infinitely Divisible Distributions*, Cambridge University Press, 1999.
- [39] R. Schilling, T. Uemura, *Dirichlet forms generated by pseudo differential operators: on the Feller property of the associated stochastic process*, to appear in Tohoku Math. J.
- [40] R. Song, Z. Vondraček, *Harnack inequality for some classes of Markov processes*, Math. Z. 246 (2004), no. 1-2, 177-202.
- [41] P. Sztonyk, *On harmonic measures for Lévy processes*, Prob. Math. Statist. 20(2) (2000), 383-390.
- [42] S. J. Taylor, *Sample path properties of a transient stable process*, J. Math. Mech. 16 (1967), 1229-1246.
- [43] H. Triebel, *Theory of function spaces*, Monographs in Mathematics, Vol. 78. Basel-Boston-Stuttgart: Birkhäuser Verlag, (1983).

- [44] T. Watanabe *Asymptotic estimates of multi-dimensional stable densities and their applications*, Trans. Amer. Math. Soc. 359 (2007), 2851-2879.
- [45] J.-M. Wu *Harmonic measures for symmetric stable processes*, Studia Math. 149 (2002), no. 3, 281–293.